

When planning to survive goes wrong: predicting the future and replaying the past in anxiety and PTSD

Christopher Gagne¹, Peter Dayan³ and Sonia J Bishop^{1,2}



We increase our probability of survival and wellbeing by minimizing our exposure to rare, extremely negative events. In this article, we examine the computations used to predict and avoid such events and to update our models of the world and action policies after their occurrence. We also consider how these computations might go wrong in anxiety disorders and Post Traumatic Stress Disorder (PTSD). We review evidence that anxiety is linked to increased simulations of the future occurrence of high cost negative events and to elevated estimates of the probability of occurrence of such events. We also review psychological theories of PTSD in the light of newer, computational models of updating through replay and simulation. We consider whether pathological levels of re-experiencing symptomatology might reflect problems reconciling the traumatic outcome with overly optimistic priors or difficulties terminating off-line simulation focused on negative events and over-generalization to states sharing features with those antecedent to the trauma.

Addresses

¹ Department of Psychology, University of California, Berkeley, CA 94720, USA

² Helen Wills Neuroscience Institute, University of California, Berkeley, CA 94720, USA

³ Gatsby Computational Neuroscience Unit, University College London, London W1T 4JG, UK

Corresponding author: Bishop, Sonia J (sbishop@berkeley.edu)

Current Opinion in Behavioral Sciences 2018, 24:89–95

This review comes from a themed issue on **Survival circuits**

Edited by **Dean Mobbs** and **Joe LeDoux**

<https://doi.org/10.1016/j.cobeha.2018.03.013>

2352-1546/© 2018 Published by Elsevier Ltd.

Introduction

In modern life, aversive events vary both in their frequency and severity. Shootings, terrorist incidents and plane crashes are rare, extremely negative events that might threaten our survival if experienced even just once. Avoiding exposure to such events and handling them appropriately if they occur is critical to our survival and well-being but, we argue, surprisingly hard to integrate smoothly into the course of our day-to-day lives. Here, we

lay out this computational problem as a form of approximate Bayesian decision theory (BDT) [1] and consider how miscalibrated attempts to solve it might contribute to anxiety and stress disorders.

According to BDT, we should combine a probability distribution over all relevant states of the world with estimates of the benefits or costs of outcomes associated with each state. We must then calculate the course of action that delivers the largest long-run expected value. Individuals can only possibly approximately solve this problem. To do so, they bring to bear different sources of information (e.g. priors, evidence, models of the world) and apply different methods to calculate the expected long-run values of alternate courses of action. Avoiding catastrophic events poses unique difficulties above those in other situations framed in BDT because the rarity of these events renders methods that work well in more typical situations, such as model-free learning, relatively less useful. As a result, model-based processes that more efficiently re-use and extend experience, such as replay and counterfactual simulation, become especially important both before and after these events.

In part 1, we discuss the computations required to take into account the potential future occurrence of yet-to-happen rare, extremely negative events as we plan and navigate our daily lives. We consider how individual differences in these computations might confer vulnerability to anxiety. In part 2, we focus on the computations required to update our models of the world and action policies after the occurrence of rare, extremely negative events. We explore how the re-experiencing symptomatology characteristic of Post Traumatic Stress Disorder (PTSD) might be understood in the context of these computations.

Part 1: anxiety and predicting the future

The survival circuits that are the focus of this issue provide rich, hard-wired (sometimes called Pavlovian) policies that directly determine particular actions in the face of immediate mortal threat. However, waiting until threats materialize is rarely wise; maximizing our chances of survival and well-being requires estimating the probability and cost of extreme negative events and developing strategies for ameliorating or avoiding them ahead of time.

When estimating the expected value of avoidance behaviors, one should weight outcome costs by outcome

probabilities. In the case of rare, extremely negative events, the costs are so high that even small differences in probability estimates will have a huge impact on these expected values. High trait anxious children and adults produce higher estimates of the probability that future negative events (e.g. being involved in a road accident) will occur to them than do low trait anxious participants [2–4]. Such differences in probability estimates might result in increased selection of avoidance behaviors despite the associated disruption to everyday life activities.

There are a number of potential sources of these anxiety-related differences in probability estimates; these include differences in the method of estimation used, in initial biases in the estimates (priors) and in the precision of estimate calculation. The probability of rare, extremely negative events is hard to calculate precisely because similar events have rarely, if ever, been actually experienced. Thus, probability estimates are likely to be broad, with weak upper bounds. If the world is rapidly changing, that is, volatile, these bounds should be weaker still, as only very recent outcomes are pertinent [5]. In anxiety, there is evidence for difficulties in estimating environmental volatility [6*,7] and increased adoption of high volatility priors [8*]. Hence, anxious individuals might have even weaker upper bounds than other individuals for probability estimates of rare, extremely negative events. One strategy for robustly avoiding catastrophic failures is to adopt a worse-case scenario (H_∞ control) [9], that is, to rely on the upper bound as opposed to the mean or median of the distribution of outcome probabilities. Consistent with this, clinically anxious individuals are reported to engage in catastrophizing, focusing on worst case outcomes [10]. If anxious individuals do indeed show a combination of widened bounds for probability estimates of rare, extremely negative events and reliance on upper bounds during action selection, this might promote more frequent selection of avoidance behaviors.

If we seek to weigh up the benefits of certain behaviors (e.g. going on vacation in London) against the potential probability and cost of rare, extremely negative events (e.g. a plane crash or terrorist incident), we must calculate the long-run values of alternate actions. Long-run values take into account outcomes that might only arise several steps after the initial choice. The methods used for this are often conceived as living on a spectrum between so-called model-free and model-based computations [11,12]. Model-free and model-based methods both aim to produce appropriate policies (which specify the actions that should be taken in different situations), however, they use information from the world differently to do so. Model-free methods such as temporal difference learning [13,14] cache information during experience of the environment. They thereby create policies that are computationally straightforward to use to guide subsequent on-the-spot

action selection and have the speed to be well suited to the avoidance or mitigation of extremely negative events. However, use of model-free methods to create multi-step action policies aimed at avoidance of rare, extremely negative events is heavily compromised by the reliance of these methods on past experience, given the typical absence of past experience for such events.

In contrast, model-based methods construct internal representations of alternate states of the world, and of how alternate future courses of action might play out depending on the initial state encountered [14,15]. Construction of such models of the world is informed by direct experience. However, indirect evidence such as vicarious experience or intuitive knowledge about the physical world can also be incorporated [16]. Sampling from the model can be used to play out what might transpire given selection of a particular initial action, even if that action has not been taken in real life. Thus, model-based methods can anticipate states and outcomes that have never been experienced, a characteristic of particular value for working out how to avoid rare, negative events. Such sampling can be used directly for planning [17]. However, it has also been suggested that sampling during off-line periods (such as quiet wakefulness or sleep) can be used to train model-free estimates of action values [18]. The putative benefit of this is to create model-free action policies that are fast to use but nevertheless reflect the knowledge contained within the model. However, if the model, or samples drawn from it, is biased, then not only will model-based planning be biased, but the model-free policy trained on the basis of the samples drawn will become biased too.

Biased sampling is likely to impact all of us, to some extent, but might be a particular problem for individuals at risk for anxiety disorders. Given the impossibility of exploring all potential future states, we need strategies for restricting the states we consider. It has been suggested that we focus on states that are easily *available* [19]. The frequency with which states have been encountered in the past is likely to impact their availability. This may result in low frequency outcomes being overlooked during the estimation of action values. However, relying on state frequency alone might be suboptimal in some situations, and it has been suggested that this might be offset by the oversampling of emotionally salient outcomes, especially those involving extreme (i.e. rare, high value) events [20**]. In line with this, emotional salience has been shown to facilitate recall of past events [21], in particular in the case of extreme events such as terrorist attacks [22,23]. That increased availability of such events might impact action valuation is indicated by findings that availability of positive and negative events during simulation and recall predicts estimates of the future probability of these events [24,25]. A current example is the reported drop in Southwest bookings the week after an engine

broke apart resulting in the death of a passenger on one of their flights.

Anxious individuals potentially oversample negative extreme outcomes and associated antecedents to a greater extent than individuals low in anxiety. In line with this, participants with high anxiety levels selectively generate more negative possible future life events than low anxious participants within a limited period of time [26]. If anxiety is linked to oversampling of negative outcomes and their antecedents, the frequency of such simulation might also moderate the extent to which estimates of the values of avoidance behaviors are influenced by sampling biases. Worry, repetitive thinking focused on future potential threat, imagined catastrophes and their possible prevention [27], is a common form of simulation of the real world. Elevated levels of worry are a defining feature of Generalized Anxiety Disorder and also characterize other anxiety disorders [28]. Anxious adults report more worry episodes and greater overall time engaged in worry [29], and anxious children report being unable to stop worrying until the focus of worry is removed [30].

Frequent, uncontrollable simulation of negative outcomes and their antecedents might contribute to the maintenance of anxiety disorders by increasing the subjective valuation, and selection, of avoidance behaviors. If anxious participants also show increased reliance on upper bounds of probability estimates for rare negative events (as discussed above), this will have a converging influence upon the overvaluation of avoidance behaviors. These behaviors, in turn, will reduce opportunities for anxious individuals to collect data showing that extreme negative events almost never occur, even if avoidance behaviors are not engaged. Therefore, there will not be the observations needed to correct estimation biases and stabilize a potentially detrimental cycle of increasing miscalibration of action value estimates and selection of avoidance behaviors. Such decision-theoretic path dependencies have been implicated in various other psychiatric contexts [31,32].

Part 2. PTSD and replaying the past

Despite our best efforts, extremely negative events do occasionally occur. If such an event is survived, the balance of planning activities should shift toward avoiding the event being experienced again. This is both because the events occurrence might contain information useful for avoiding similar events in the future and because there might be autocorrelation in the occurrence of extremely negative events (e.g. when new predators enter an environment [33]).

Off-line replay of prior experiences and simulation of counterfactual actions and associated outcomes provide a means to update action values following the occurrence of an extremely negative event [18]. It has been argued that


previous states should be prioritized for replay based on how much that replay would change value estimates [34**]. One way to accomplish this is by tagging states based on how much their successors value has changed [35]. If change in value estimates determines priority for replay, the astronomically large discrepancy in outcome value occasioned by the occurrence of a rare, extremely negative event would be expected to result in prioritized replay of that events antecedents (see Figure 1).

By replaying the states that preceded a rare, extremely negative event, we can ascribe more negative values to these states and the actions selected within them that led up to the events occurrence. Equally, actions that were not taken can be simulated, together with the possible outcomes of these actions. Should similar states be encountered again, the model-free system can use the updated action values to choose swiftly a safer course of action. However, this may not be entirely straightforward. Specifically, exposure to a rare, extreme event might increase the salience and availability of similar outcomes and increase the probability, from an otherwise negligible level, of such outcomes being simulated following various actions. This might result in many courses of action being evaluated more negatively than before the experience of the extreme negative event.

Findings from the traumatic stress literature indicate that most individuals do indeed replay the antecedents of extreme negative events after their occurrence. Following extreme negative (also termed 'traumatic') events, such as motor-vehicle accidents, over 50% of individuals report intrusive recollections, flashbacks, or nightmares up to three months following the event [36,37]. These phenomena are collectively referred to as re-experiencing symptomatology. It is also common for people to ruminate repeatedly on their experience, thinking about the events causes and ways that it might have been prevented, for example, 'I was running late so I cut through town, if I had gone the long way round . . .' [36,38,39].

Some degree of re-experiencing, rumination and counterfactual thinking about the past might be functional in the wake of an extremely negative event, or trauma. Indeed, psychological accounts have argued that repetitive thinking in the wake of a traumatic event might be important for resolving the discrepancy between the event and pre-existing core beliefs or assumptions. Horowitz [40] describes this process as 'cognitive processing' and Janoff-Bulman [41] describes it as 'integration', resulting in what Tedeschi and Calhoun [42] describe as 'post-traumatic growth'. Critically, whereas re-experiencing symptoms decay rapidly over time for some individuals, for others they remain frequent and cause significant levels of distress and disruption to everyday life. Researchers have struggled to identify aspects of the re-experiencing process or content that predict

Figure 1



A replay and simulation account of PTSD. When a traumatic event is experienced, the difference in value between the expected and actual outcome is calculated (box 2). Given the high negative value of the traumatic event and its low prior probability of occurrence, a large negative prediction error (δ) will be experienced. This prediction error triggers the replay of the antecedent state and action (box 3). During this replay, the antecedent state and action will be ascribed a more negative value. Alternate actions will also be simulated in order to identify a better counterfactual course of action from the antecedent state. The action values employed when the initial decision was made (box 1) will be updated. Pre-trauma, the sampling, and hence consideration during action valuation, of traumatic outcomes is unlikely. Post-trauma, the increased salience and availability of such outcomes will make their sampling more probable. Even very small increases in the estimated probability of actions leading to the same, or other, traumatic outcomes will result in substantial negative revision of action values given the large negative value of such outcomes. Replay and simulation will continue until a counterfactual action is found with a sufficiently large expected value ($Q(S_t, a)$) or until the prediction errors (δ) resulting from the replay and simulations have sufficiently diminished. Increased estimates of the probability of traumatic outcomes and associated downward revision of action values may result in many actions being considered in search for one with an acceptable value. Stopping criteria (thresholds for $Q(S_t, a)$ or δ) may vary across individuals and may potentially be influenced by external factors (e.g. novel stressors). Other states that share perceptual or semantic features with the state antecedent to the traumatic event will likely also be replayed or simulated and their associated actions may be ascribed a more negative value to the extent that shared features increase the availability, during simulation, of traumatic outcomes (box 4). The change in value of the state immediately antecedent to the traumatic event (S_t) will entail that the values of states and actions (starting from S_{t-1}, A_{t-1}) before that state will also need to be updated [34^a,35]. This will result in the replay and re-valuation of increasingly earlier states and actions (box 5). Generalization of values to similar states and the simulation and re-valuation of alternate courses of action will also occur from these more distant antecedents. This cycle of replay will continue, with gradual discounting as more temporally distant states are revisited, until prediction errors (δ) sufficiently diminish or until counterfactual actions can be found at each temporal point in the antecedent chain and at each level of generalization with sufficiently high expected values. This process of replay of chosen action paths and simulation of alternate action paths might be phenomenologically experienced as intrusive thoughts, dreams, rumination and counterfactual reasoning. Red text is used to signify points where individual differences might confer vulnerability to elevated PTSD symptomatology. (a) Individuals may differ in stopping criteria; for example, individuals vulnerable to PTSD might be reluctant to take actions with even the slightest possibility of future catastrophe or might have a lower tolerance for negative changes in action or state values. (b) A reduced rate of discounting of the prediction error as more temporally distant antecedents are considered and (c) a shallower generalization slope when re-valuing states or actions that resemble those antecedent to the trauma are likely to increase the number of states and actions replayed resulting in higher levels of re-experiencing, avoidance and hyper-vigilance.

post-traumatic growth versus disorder [43]. Here, we suggest that the re-experiencing symptomatology and negative cognitions (e.g. rumination and counterfactual reasoning) observed following an extreme negative event can be computationally operationalized in terms of replay and simulation. In the rest of this section, we consider how this operationalization might shed light on the potential determinants of healthy versus dis-ordered responses to the experience of an extremely negative event.

Several psychological accounts of PTSD posit that individuals with rigid beliefs about the positive nature of the world are more likely to experience PTSD after a

traumatic event due to their assumptions or schema about the world being unable to flexibly accommodate or integrate the traumatic experience [41,44]. According to BDT, if an individual has a much lower prior expectation of the occurrence of extremely negative events, this will generate a larger discrepancy between the value of the expected outcome and the value of the actual outcome (the traumatic event), strongly prioritizing the replay of the events antecedent states and actions [34^a,35] and likely resulting in greater re-experiencing symptomatology. If rigidity is associated with an unwillingness to update state and action values to make them more negative, then negative prediction errors will stubbornly persist.

Individuals who go on to develop PTSD endorse more negative world views in the immediate aftermath of trauma [45]. Moreover, elevated pre-trauma levels of anxiety and depression have been found prospectively to predict levels of post-traumatic stress symptomatology [46]. Since anxiety and depression are linked to negative, not positive, biases in beliefs, interpretations of ambiguous events and judgements about the future [47–49], it seems unlikely that the magnitude of the prediction error occasioned by experiencing a traumatic event is a key predisposing factor, at least for these individuals. Indeed, there is little empirical evidence that pre-trauma possession of optimistic priors confers vulnerability to PTSD.

There are other ways in which individual differences might influence extent of engagement in replay and simulation. One possibility is that individuals with a pre-trauma history of anxiety or depression might be more prone to difficulties with terminating disadvantageous replay or simulation processes. Such stopping difficulties might confer vulnerability to PTSD as well as to anxiety disorders (as described in Section 1) and depressive disorders. In line with this, both anxiety and depression are characterized by elevated levels of repetitive thoughts (worry and rumination) [50]. Further, pre-trauma engagement in repetitive thinking is a significant predictor of post-trauma levels of PTSD symptomatology [51]. In the anxiety literature, difficulty finding a potential course of action with a sufficiently positive expected value has been associated with increased uncontrollability of worry [52]. Post-trauma, inability to identify one or more counterfactual courses of action with high enough expected values to terminate simulation and replay might similarly be linked to elevated PTSD symptomatology (Figure 1).

An increased propensity for repetitive thinking might be further compounded by a disposition to over-generalization [53]. Clinical accounts of PTSD describe how everyday sounds, like a balloon popping or a car backfiring, that resemble a gunshot can lead to extreme physiological and emotional responses in individuals for whom the traumatic event involved combat or gun violence. This has led to suggestions that over-generalization might be a key vulnerability factor in PTSD [38]. In terms of the replay and simulation framework put forward here (see Figure 1), when an individual uses replay and simulation to update the value of states and actions that preceded a traumatic event, a key issue will be determining how far to go in also updating the value of other states and actions that share features with the antecedent states and actions. Other states may be related to antecedent states at a very concrete level (e.g. a similar looking street-corner to where the accident occurred) or at a very abstract level (e.g. any form of transportation in which you are not in control). Both selection of an abstract level and, at any given level, adoption of less steep (dis)similarity gradients might result in larger numbers of states and actions being

re-evaluated both off-line and on-line when the individual encounters a state that shares features with a state antecedent to the traumatic event. Future planning would suffer from a similar problem.

There is strong evidence linking both forms of over-generalization described above to anxiety, depression and PTSD. Patients with GAD, Panic Disorder, and PTSD have been shown to generalize from a conditioned stimulus across a wider range of perceptually similar shapes than healthy controls [54,55,56*]. In addition, over-general autobiographical recall has been shown to characterize both patients with depressive disorders and individuals with a history of trauma [57]. Further, levels of rumination have been shown to increase the influence of over-general memory on both future depressive and post-traumatic stress symptomatology [58,59]. Trauma-analog studies have also reported that participants asked to ruminate abstractly, versus concretely, after viewing a traumatic video show both prolonged negative mood and more negative intrusions [60,61].

In addition to re-experiencing, PTSD is also characterized by hyper-arousal and avoidance behaviors, together with other symptoms [28]. Within a replay and simulation account of PTSD, both over-generalization and going further back in the chain of antecedent states as part of the replay process (see Figure 1) will result in more states and associated actions being re-evaluated as potentially dangerous. This, in turn, could lead to an increased sense of current threat, associated physiological reactivity and avoidance of multiple situations.

One important question for future research is whether individuals with a prior history of anxiety or depression are equally likely to show over-generalization between perceptually similar stimuli or states and over-abstract levels of simulation, or if the former might be more associated with anxiety and the latter with depression. In addition, the extent to which individuals vary in stopping criteria for terminating replay and simulation, and the role of this in anxiety, depression and PTSD, remains to be established.

Conclusion

In order to survive and maximize our wellbeing, we need to weigh up actions aimed at avoiding life threatening events against the pursuit of rewarding activities and avoidance of more minor aversive outcomes or losses. Here, we have outlined the computational processes involved in action valuation both in advance of, and subsequent to, the occurrence of rare, extremely negative events, and discussed how both anxiety and PTSD might be understood within this computational framework. Future work would valuably extend our computational analysis to consideration of potential neurobiological markers of disease risk. Given the putative role of hippocampal function in off-line

model-based simulation processes [62–64] and evidence for hippocampal dysfunction in both anxiety [65–68] and PTSD [69–71], this is a particular structure of interest.

Conflict of interest statement

Nothing declared.

Acknowledgement

PD is currently on a leave of absence at Uber Technologies.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Berger JO: *Statistical Decision Theory and Bayesian Analysis*. New York: Springer; 1985.
2. Butler G, Mathews A: **Cognitive processes in anxiety**. *Adv Behav Res Ther* 1983, **5**:51–62.
3. MacLeod AK, Byrne A, Valentine JD: **Affect, emotional disorder and future-directed thinking**. *Cogn Emot* 1996, **10**:69–86.
4. Muris P, van der Heiden S: **Anxiety, depression, and judgments about the probability of future negative and positive events in children**. *J Anxiety Disord* 2006, **20**:252–261.
5. Behrens TEJ, Woolrich MW, Walton ME, Rushworth MFS: **Learning the value of information in an uncertain world**. *Nat Neurosci* 2007, **10**:1214–1221.
6. Browning M, Behrens TE, Jocham G, O'reilly JX, Bishop SJ: **Anxious individuals have difficulty learning the causal statistics of aversive environments**. *Nat Neurosci* 2015, **18**:590.
Human subjects chose between two shapes, aiming to avoid electric shocks of varying magnitudes. Volatility was varied within the task; in one phase, the shape with a higher probability of leading to shock was fixed, in the other phase, it varied across trials. Modeling of choice behavior and pupilometry revealed that high anxious subjects showed poorer tracking of volatility and reduced adjustment of learning rate between task phases.
7. Pulcu E, Browning M: **Correction: affective bias as a rational response to the statistics of rewards and punishments**. *eLife* 2017, **6**:e32902.
8. Huang H, Thompson W, Paulus MP: **Computational dysfunctions in anxiety: failure to differentiate signal from noise**. *Biol Psychiatry* 2017, **82**:440–446.
Human subjects had to locate a given target (a rightward or leftward moving random-dot stimulus), which would regularly switch among three alternate locations. Anxiety was linked to increased post-error switching; this corresponded to a higher base learning rate. This finding might indicate that anxious individuals adopt a high volatility prior when it is difficult to track the statistical regularities of a noisy signal.
9. Doyle JC, Glover K, Khargonekar PP, Francis BA: **State-space solutions to standard H_2 and H_∞ control problems**. *IEEE Trans Autom Control* 1989, **34**:831–847.
10. Sandin B, Sánchez-Arribas C, Chorot P, Valiente RM: **Anxiety sensitivity, catastrophic misinterpretations and panic self-efficacy in the prediction of panic disorder severity: towards a tripartite cognitive model of panic disorder**. *Behav Res Ther* 2015, **67**:30–40.
11. Doya K: **What are the computations of the cerebellum, the basal ganglia and the cerebral cortex?** *Neural Netw* 1999, **12**:961–974.
12. Daw ND, Niv Y, Dayan P: **Uncertainty-based competition between prefrontal and dorsolateral striatal systems for behavioral control**. *Nat Neurosci* 2005, **8**:1704.
13. Sutton RS: **Learning to predict by the methods of temporal differences**. *Mach Learn* 1988, **3**:9–44.
14. Sutton RS, Barto AG: *Reinforcement Learning: An Introduction*. Cambridge: MIT Press; 1998.
15. Daw ND, Dayan P: **The algorithmic anatomy of model-based evaluation**. *Philos Trans R Soc B* 2014, **369**:20130478.
16. Battaglia PW, Hamrick JB, Tenenbaum JB: **Simulation as an engine of physical scene understanding**. *Proc Natl Acad Sci U S A* 2013, **110**:18327–18332.
17. Kocsis L, Szepesvári C: **Bandit based Monte-Carlo planning**. In *Machine Learning: ECML 2006*. Edited by Fürnkranz J, Scheffer T, Spiliopoulou M. Berlin Heidelberg: Springer; 2006:282–293.
18. Sutton RS: **Integrated architectures for learning, planning, and reacting based on approximating dynamic programming**. In *Machine Learning Proceedings 1990*. Elsevier; 1990:216–224.
19. Kahneman D, Slovic P, Tversky A: **The simulation heuristic**. *Judgment Under Uncertainty: Heuristics and Biases*. Cambridge University Press; 1982.
20. Lieder F, Griffiths TL, Hsu M: **Overrepresentation of extreme events in decision making reflects rational use of cognitive resources**. *Psychol Rev* 2018, **125**:1.
This paper considers how we sample from memory when making decisions involving extreme (i.e. rare, high value) negative or positive outcomes. According to the account put forward, sampling low frequency outcomes that are readily available as a result of their emotional salience is often optimal given cognitive constraints. The effect of sampling bias on estimation error is held to be offset by a reduction in sampling variance.
21. Dolcos F, Katsumi Y, Weymar M, Moore M, Tsukiura T, Dolcos S: **Emerging directions in emotional episodic memory**. *Front Psychol* 2017, **8**.
22. Lichtenstein S, Slovic P, Fischhoff B, Layman M, Combs B: **Judged frequency of lethal events**. *J Exp Psychol [Hum Learn]* 1978, **4**:551.
23. Madan CR, Ludvig EA, Spetch ML: **Remembering the best and worst of times: memories for extreme outcomes bias risky decisions**. *Psychon Bull Rev* 2014, **21**:629–636.
24. MacLeod AK, Williams JM, Bekerian DA: **Worry is reasonable: the role of explanations in pessimism about future personal events**. *J Abnorm Psychol* 1991, **100**:478.
25. MacLeod C, Campbell L: **Memory accessibility and probability judgments: an experimental evaluation of the availability heuristic**. *J Pers Soc Psychol* 1992, **63**:890.
26. MacLeod AK, Byrne A: **Anxiety, depression, and the anticipation of future positive and negative experiences**. *J Abnorm Psychol* 1996, **105**:286.
27. Watkins ER: **Constructive and unconstructive repetitive thought**. *Psychol Bull* 2008, **134**:163.
28. AP Association: *Diagnostic and Statistical Manual of Mental Disorders (DSM-5®)*. American Psychiatric Pub; 2013.
29. Verkuil B, Brosschot JF, Thayer JF: **Capturing worry in daily life: are trait questionnaires sufficient?** *Behav Res Ther* 2007, **45**:1835–1844.
30. Szabó M, Lovibond PF: **The cognitive content of thought-listed worry episodes in clinic-referred anxious and nonreferred children**. *J Clin Child Adolesc Psychol* 2004, **33**:613–622.
31. Huys QJ, Daw ND, Dayan P: **Depression: a decision-theoretic analysis**. *Annu Rev Neurosci* 2015, **38**:1–23.
32. Dayan P, Roiser JP, Viding E: **The first steps on long marches: the costs of active observation**. In *Rethinking Biopsychosocial Psychiatry*. Edited by Davies W, Roache R, Savulescu J. Oxford University Press; 2018. (in press).
33. Travis JM, Palmer SC, Coyne S, Millon A, Lambin X: **Evolution of predator dispersal in relation to spatio-temporal prey dynamics: how not to get stuck in the wrong place!** *PLOS ONE* 2013, **8**:e54453.
34. Mattar MG, Daw ND: **Prioritized memory access explains planning and hippocampal replay**. *bioRxiv* 2018.
Presents a theoretical account for how the states replayed during policy updating are prioritized. The theory proposes that priority for replay

should be given to states that have the largest potential to change expected utility if included in the policy update. Under this theory, prioritized states are often those antecedent to outcomes that generate the largest prediction errors. The theory accommodates findings from the hippocampal replay literature, such as the change in reverse-replay frequency following increases or decreases in reward.

35. Moore AW, Atkeson CG: **Prioritized sweeping: reinforcement learning with less data and less time.** *Mach Learn* 1993, **13**:103-130.
 36. Ehlers A, Mayou RA, Bryant B: **Psychological predictors of chronic posttraumatic stress disorder after motor vehicle accidents.** *J Abnorm Psychol* 1998, **107**:508.
 37. Bryant R: **Acute stress disorder.** *PTSD Res Q* 2000, **11**.
 38. Ehlers A, Clark DM: **A cognitive model of posttraumatic stress disorder.** *Behav Res Ther* 2000, **38**:319-345.
 39. El Leithy S, Brown GP, Robbins I: **Counterfactual thinking and posttraumatic stress reactions.** *J Abnorm Psychol* 2006, **115**:629.
 40. Horowitz MJ: **Disasters and psychological responses to stress.** *Psychiatr Ann* 1985, **15**:161-167.
 41. Janoff-Bulman R: *Shattered Assumptions: Towards a New Psychology of Trauma.* Free Press; 1992.
 42. Tedeschi RG, Calhoun LG: **Posttraumatic growth: conceptual foundations and empirical evidence.** *Psychol Inq* 2004, **15**:1-18.
 43. Dekel S, Mandl C, Solomon Z: **Shared and unique predictors of post-traumatic growth and distress.** *J Clin Psychol* 2011, **67**:241-252.
 44. Park CL, Mills MA, Edmondson D: **PTSD as meaning violation: testing a cognitive worldview perspective.** *Psychol Trauma Theory Res Pract Policy* 2012, **4**:66.
 45. Ginzburg K: **PTSD and world assumptions following myocardial infarction: a longitudinal study.** *Am J Orthopsychiatry* 2004, **74**:286.
 46. Orr SP, Lasko NB, Macklin ML, Pineles SL, Chang Y, Pitman RK: **Predicting post-trauma stress symptoms from pre-trauma psychophysiological reactivity, personality traits and measures of psychopathology.** *Biol Mood Anxiety Disord* 2012, **2**:8.
 47. Beck AT: *Cognitive Therapy and the Emotional Disorders.* International Universities Press; 1976.
 48. Teasdale JD: **Negative thinking in depression: cause, effect, or reciprocal relationship?** *Adv Behav Res Ther* 1983, **5**:3-25.
 49. Mathews A, MacLeod C: **Cognitive vulnerability to emotional disorders.** *Annu Rev Clin Psychol* 2005, **1**:167-195.
 50. Kircanski K, Thompson RJ, Sorenson JE, Sherdell L, Gotlib IH: **Rumination and worry in daily life: examining the naturalistic validity of theoretical constructs.** *Clin Psychol Sci* 2015, **3**:926-939.
 51. Spinhoven P, Penninx BW, Krempeuiou A, van Hemert AM, Elzinga B: **Trait rumination predicts onset of Post-Traumatic Stress Disorder through trauma-related cognitive appraisals: a 4-year longitudinal study.** *Behav Res Ther* 2015, **71**:101-109.
 52. Szabó M, Lovibond PF: **Worry episodes and perceived problem solving: a diary-based approach.** *Anxiety Stress Coping* 2006, **19**:175-187.
 53. Watkins E: **Dysregulation in level of goal and action identification across psychological disorders.** *Clin Psychol Rev* 2011, **31**:260-278.
 54. Lissek S: **Toward an account of clinical anxiety predicated on basic, neurally mapped mechanisms of Pavlovian fear-learning: the case for conditioned overgeneralization.** *Depress Anxiety* 2012, **29**:257-263.
 55. Lissek S, Kaczurkin AN, Rabin S, Geraci M, Pine DS, Grillon C: **Generalized anxiety disorder is associated with overgeneralization of classically conditioned fear.** *Biol Psychiatry* 2014, **75**:909-915.
 56. Kaczurkin AN, Burton PC, Chazin SM, Manbeck AB, Espensen-Sturges T, Cooper SE, Sponheim SR, Lissek S: **Neural substrates of overgeneralized conditioned fear in PTSD.** *Am J Psychiatry* 2016, **174**:125-134.
- Rings varying in size were presented to patients with PTSD; either the smallest or largest ring was paired with a shock. Relative to controls, patients with PTSD had shallower generalization gradients, meaning that they showed a more gradual drop off in conditioned fear responses as rings deviated further in size from the one paired with shock.
57. Williams JMG, Barnhofer T, Crane C, Herman D, Raes F, Watkins E, Dalgleish T: **Autobiographical memory specificity and emotional disorder.** *Psychol Bull* 2007, **133**:122.
 58. Hamlat EJ, Connolly SL, Hamilton JL, Stange JP, Abramson LY, Alloy LB: **Rumination and overgeneral autobiographical memory in adolescents: an integration of cognitive vulnerabilities to depression.** *J Youth Adolesc* 2015, **44**:806-818.
 59. Kleim B, Ehlers A: **Reduced autobiographical memory specificity predicts depression and posttraumatic stress disorder after recent trauma.** *J Consult Clin Psychol* 2008, **76**:231.
 60. Ehring T, Frank S, Ehlers A: **The role of rumination and reduced concreteness in the maintenance of posttraumatic stress disorder and depression following trauma.** *Cogn Ther Res* 2008, **32**:488-506.
 61. Ehring T, Fuchs N, Kläsener I: **The effects of experimentally induced rumination versus distraction on analogue posttraumatic stress symptoms.** *Behav Ther* 2009, **40**:403-413.
 62. Wilson MA, McNaughton BL: **Reactivation of hippocampal ensemble memories during sleep.** *Science* 1994, **265**:676-679.
 63. Foster DJ, Wilson MA: **Reverse replay of behavioural sequences in hippocampal place cells during the awake state.** *Nature* 2006, **440**:680.
 64. Ambrose RE, Pfeiffer BE, Foster DJ: **Reverse replay of hippocampal place cells is uniquely modulated by changing reward.** *Neuron* 2016, **91**:1124-1136.
 65. Gray JA, McNaughton N: *The Neuropsychology of Anxiety: An Enquiry into the Function of the Septo-Hippocampal System.* Oxford University Press; 2003.
 66. Khemka S, Barnes G, Dolan RJ, Bach DR: **Dissecting the function of hippocampal oscillations in a human anxiety model.** *J Neurosci* 2017, **37**:6869-6876.
 67. Bach DR, Guitart-Masip M, Packard PA, Miró J, Falip M, Fuentemilla L, Dolan RJ: **Human hippocampus arbitrates approach-avoidance conflict.** *Curr Biol* 2014, **24**:541-547.
 68. Oler JA, Fox AS, Shelton SE, Rogers J, Dyer TD, Davidson RJ, Shelledy W, Oakes TR, Blangero J, Kalin NH: **Amygdalar and hippocampal substrates of anxious temperament differ in their heritability.** *Nature* 2010, **466**:864.
 69. Shin LM, Rauch SL, Pitman RK: **Amygdala, medial prefrontal cortex, and hippocampal function in PTSD.** *Ann N Y Acad Sci* 2006, **1071**:67-79.
 70. Kheirbek MA, Klemenhagen KC, Sahay A, Hen R: **Neurogenesis and generalization: a new approach to stratify and treat anxiety disorders.** *Nat Neurosci* 2012, **15**:1613.
 71. Stevens JS, Reddy R, Kim YJ, van Rooij SJ, Ely TD, Hamann S, Ressler KJ, Jovanovic T: **Episodic memory after trauma exposure: medial temporal lobe function is positively related to re-experiencing and inversely related to negative affect symptoms.** *NeuroImage Clin* 2018, **17**:650-658.